

Identifying risk factors for bladder catheterization in Internal Medicine patients: can it be helpful for clinical practice?

Maria João Lobão^{1,2}, Paulo Sousa²

¹ Internal Medicine Department, Hospital de Cascais, ² National School of Public Health, NOVA University of Lisbon

Background

Urinary tract infections are one of the most common nosocomial infections worldwide and the vast majority of it is indwelling catheter related.¹ Bladder catheterization is a very frequent procedure and it should be done strictly accomplishing the indications for it.¹ In many cases, catheters are placed inappropriately leading to unnecessary and prolonged use which can result in avoidable infections.¹⁻³ The morbidity, mortality and costs associated to catheter-associated urinary tract infections (CAUTI) are not negligible.^{4,5}

With this study, we aim to identify risk factors for bladder catheterization in patients admitted to an internal medicine ward that could help clinical reasoning and decision about catheter placement in addition to formal recommended indications.

Methods

We performed a historical cohort study that included a systematic random sample of 388 patients, representative of the 3492 admissions occurred in a Portuguese internal medicine ward (93-beds) in 2014. Patients transferred from or to another hospital, as well as patients with admission diagnosis of urinary tract infection were excluded. Variables related to patient (age, sex, age adjusted Charlson comorbidity index^{6,7}, place of residency, functional⁸ and nutritional status⁹, sphincter incontinence, pressure ulcers), and to admission episode (department from admission, principal diagnosis and length of stay) were analyzed.

Univariate analysis was done to characterize the cohort. We performed a bivariate analysis to identify statistically significant variables that were associated to bladder catheterization. Binary Logistic Regression (enter method) was used to identify independent risk factors for catheter use and to develop the predictive risk model. *Odds Ratio* and 95% confidence intervals were calculated. We tested several consecutive models until the final one that only considered variables with p-value < 0.05 (value considered to be statistically significant in this study). We performed Hosmer and Lemeshow to test the goodness of fit of the model. The Receiver Operating Characteristic (ROC) curve was used as a measure of the model's predictive discrimination.¹⁰ Statistical analysis was performed with SPSS IBM® Statistics Software (24th version).

Results

Figure 1 Cohort patients inclusion diagram





Figure 2 Cohort characterization and main results



Bivariate analysis

Variables related to the Patient			Bladder Catheterization (n=96)	No Bladder Catheterization (n=292)	Relative Risk (CI 95%)	Odds Ratio (CI 95%)	p-value
Gender	woman	n (%)	56 (58.3%)	143 (71.9%)	1.32 (0.59-1.89)	1.45 (0.91-2.30)	p=0.111 [§]
	man	n (%)	40 (41.7%)	149 (51%)	1.0	1.0	
Age	< 65 Years	n (%)	7 (7.3%)	93 (31.8%)	0.22 (0.10-0.47)	0.16 (0.07-0.37)	p<0.001 [§]
	65 or > 65 Years	n (%)	89 (92.7%)	199 (68.2%)	1.0	1.0	
Place of residence	Own home	n (%)	71 (74.0%)	259 (88.7%)	0.49 (0.34-0.71)	0.36 (0.20-0.64)	p<0.001 [§]
	Institution	n (%)	25 (26.9%)	33 (11.3%)	1.0	1.0	
Diabetes Mellitus	Presence	n (%)	22 (22.9%)	53 (18.2%)	1.24 (0.62-1.85)	1.34 (0.76-2.3)	p=0.305 [§]
	Absence	n (%)	74 (77.1%)	239 (81.8%)	1.0	1.0	
Cancer	Presence	n (%)	14 (14.6%)	51 (17.5%)	0.84 (0.5-1.39)	0.86 (0.42-1.53)	p=0.512 [§]
	Absence	n (%)	82 (85.4%)	241 (82.5%)	1.0	1.0	
Immunosuppression	Presence	n (%)	8 (8.3%)	27 (9.2%)	0.91 (0.48-1.73)	0.89 (0.39-2.03)	p=0.786 [§]
	Absence	n (%)	88 (91.7%)	265 (90.8%)	1.0	1.0	
Age adjusted Charlson Comorbidity Index	average (SD)		5.05 (±1.61)	3.53 (±2.23)	n.a	n.a	p<0.001 [§]
	median (IQR)		5 (4-6)	4 (3-5)	n.a	n.a	
Nutritional Status	High Risk	n (%)	35 (25.2%)	67 (22.9%)	1.68 (1.13-2.28)	1.92 (1.17-3.16)	p=0.009 [§]
	Low Risk	n (%)	61 (70.8%)	225 (77.1%)	1.0	1.0	
Sphincter incontinence	Presence	n (%)	83 (86.5%)	121 (41.4%)	5.75 (3.32-9.97)	9.02 (4.8-16.92)	p<0.001 [§]
	Absence	n (%)	13 (13.5%)	171 (58.6%)	1.0	1.0	
Pressure Ulcers	Presence	n (%)	28 (29.2%)	38 (13.0%)	2.0 (1.4-2.8)	2.75 (1.5-4.8)	p<0.001 [§]
	Absence	n (%)	68 (70.8%)	254 (87.0%)	1.0	1.0	

Legend: n.a – not applicable; SD – Standard deviation; IQR – Interquartile range; § – Qui-Square test; § – Mann-Whitney U Test


Variables related to admission episode			Bladder Catheterization (n=96)	No Bladder Catheterization (n=292)	Relative Risk (CI 95%)	Odds Ratio (CI 95%)	p-value
Department from admission	urgency	n (%)	82 (85.4%)	272 (93.2%)	1.04(0.32-3.51)	1.05(0.21-5.17)	p=0.6 [§]
	ICU	n (%)	12 (12.5%)	13 (4.5%)	2.16 (0.59-/82)	3.2 (0.55-18.71)	p=0.25 ^{§§}
	other	n (%)	2 (2.1%)	7 (2.4%)	1.0	1.0	
ICD-10 Principal Diagnosis	Diseases of the circulatory system	n (%)	29 (30.2%)	84 (28.8%)	0.96 (0.59-1.56)	0.94 (0.48-1.84)	p=0.88 [§]
	Diseases of the digestive system	n (%)	4 (4.2%)	35 (12.0%)	0.38 (0.14-1.04)	0.31 (0.09-0.99)	p=0.04 ^{§§}
	Diseases of the respiratory system	n (%)	39 (40.6%)	98 (33.6%)	1.05 (0.66-1.68)	1.08 (0.57-2.03)	p=0.07 [§]
	Neoplasms	n (%)	4 (4.2%)	20 (6.8%)	0.62 (0.23-1.64)	0.55(0.1-1.8)	p=0.33 ^{§§}
	Others	n (%)	20 (20.8%)	55 (18.8%)	1.0	1.0	
Length of stay		n.a	average (SD) median (IQR)	12.05 (± 6.95) 10 (7.0-10.0)	8.34 (± 5.06) 7 (5.0-11.0)	n.a n.a	p<0.001 ^{§§}
Barthel Index for activities of daily life	Totally dependent	n (%)	54(56.3%)	58(19.9%)	24.3 (6.09-97.31)	46.1(10.83-196.09)	p<0.001 ^{§§}
	very dependent	n (%)	36 (37.5%)	82 (28.1%)	15.3 (3.77-61.68)	21.8 (3.77-61.68)	p<0.001 ^{§§}
	Partially dependent	n (%)	4 (4.2%)	53 (18.2%)	3.5 (0.66-18.75)	3.7 (0.66-21.07)	p=0.18 ^{§§}
	Independent	n (%)	2 (2.1%)	99 (33.9%)	1.0	1.0	

Multivariate analysis and predictive risk model

Variable	OR	OR CI 95%	p [§]
Barthel Index for activities of daily life	n.a	n.a	p<0,001
Totally dependent	24,47	[5,50-108,87]	p<0,001
very dependent	11,43	[2,56-50,93]	p=0,001
Age adjusted Charlson Comorbidity Index	1,19	[1,03-1,38]	P=0,017
Length of stay	1,08	[1,04-1,13]	p<0,001

Legend: n.a – not applicable; OR – Odds ratio; OR IC95% – Confidence interval for Odds ratio; § – p-value, Wald Test
The variables with significant p-value (n=9) on the bivariate analysis were all initially introduced in the model. The variables in the final model are the ones that figures in the upper table.

Multivariate Predictive model features	
Statistic Significance	p<0,001
Hosmer e Lemeshow goodness of fit test	p=0,887



Properties of the model	
Sensitivity	34,4%
Specificity	92,5%
Positive predictive value	60,0%
Negative predictive value	81,1%
Accuracy	78,1%

Discussion and Conclusions

In this study we found a high percentage of inappropriate bladder catheterization (36,5%).^{1,11,12} This finding, that deserves further study and improve quality initiatives, underscores the need of judicious clinical reasoning about indications and benefits of this procedure for each patient individually. The independent risks factors for bladder catheterization found were total dependency and very high dependency for activities of daily life on Barthel Index on admission, Age adjusted Charlson Comorbidity Index and length of stay. The risk model can be used as a complement tool for clinical decision. Given their characteristics (high specificity and low sensibility) it only can be used to identify patients at risk for whom it must be important to establish a closer surveillance and individual decision strategy, avoiding the procedure unless it is really necessary. As we know from the literature the best way to avoid CAUTI is to restrict the use of catheters and guarantee the implementation and accomplishment of prevention bundles.^{1,3,12}

References

- Gould C V, Umscheid CA, Rajender, Agarwal K, Kurtz G, Pegues DA. Guideline for Prevention of Catheter-Associated Urinary Tract Infections (2009) GUIDELINE FOR PREVENTION OF CATHETER-ASSOCIATED URINARY TRACT INFECTIONS 2009 [Internet]. [cited 2019 Feb 15]. Available from: <https://www.cdc.gov/infectioncontrol/pdf/guidelines/cauti-guidelines.pdf>.
- Zhenoweth CE, Gould C V, Saint S. Diagnosis, management, and prevention of catheter-associated urinary tract infections [Internet]. Vol. 28, Infectious Disease Clinics of North America. 2014 [cited 2019 Feb 22]. p. 105–19. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0883552013000731>.
- Nicoll LE. Catheter-related urinary tract infection: Practical management in the elderly. Drugs and Aging. 2014 ;31(1):1–10.
- Umscheid CA, Mitchell MD, Doshi JA, Agarwal R, Williams K, Brennan PJ. Estimating the Proportion of Healthcare-Associated Infections That Are Reasonably Preventable and the Related Mortality and Costs. Infect Control Hosp Epidemiol. 2011;32(02):101–14.
- Klevers RM, Edwards JR, Richards CL, Horan TC, Gaynes RP, Pollock DA, et al. Estimating Health Care-Associated Infections and Deaths in U.S. Hospitals, 2002. Public Health Rep. 2007;122(2):160–6.
- Sundararajan V, Henderson T, Perry C, Muggivan A, Quan H, Ghali WA. New ICD-10 version of the Charlson comorbidity index predicted in-hospital mortality. J Clin Epidemiol. 2004 ;57(12):1288–94.
- Charlson ME, Pompei P, Ales KL, MacKenzie CR. A new method of classifying prognostic comorbidity in longitudinal studies: Development and validation. J Chronic Dis. 1987;40(5):373–83.
- A8. Cid-Ruza J, Damir-Moreno J. Rev Esp Salud Publica. 2009;71(2):127–37.
- Kondrup J, Allison SP, Elia M, Vellas B, Plauth M, Educational and Clinical Practice Committee ES of P and EN (ESPEN). ESPEN guidelines for nutrition screening 2002. Clin Nutr. 2003;22(4):415–21.
- Aguiar P. Estatística em Investigação Epidemiológica: SPSS Guia prático de Medicina. First. Climeps, editor; Lisboa; 2007. 237 p.
- Munasinghe RL, Yazdani H, Siddique M, Hafeez W. Appropriateness of Use of Indwelling Urinary Catheters in Patients Admitted to the Medical Service. Infect Control Hosp Epidemiol. 2001;22(10):64–12.
- Vincitorio D, Barbadore P, Pennacchietti L, Pellegrini I, David S, Ponzio E, et al. Risk factors for catheter-associated urinary tract infection in Italian elderly. Am J Infect Control. 2014;42(8):898–901.